

Secondary Hyper Parathyroidism, a Typical Mineral and Bone Disorder in a Chronic Hemodialysis Patient in Ziguinchor, Southern Senegal

Kane Yaya^{1*}, Bangoura Mohamed¹, Dia Gueye Diatou³, Aidara Chérif Mohamadou¹, Lemrabott Ahmet Tall², Faye Maria², Fall Khodia², Faye Moustapha¹, Diawara Mame Selly⁴, Diallo Penda², Cisse M Moustapha⁴, Seck S Mohamed³, Ka El Hadji Fary², Niang Abdou², Diouf Boucar²

¹Nephrology and Haemodialysis Department – Orthopedic, Radiology department, Hôpital de la paix, CHRZ, Assane Seck University, Ziguinchor, Senegal.

²Nephrology and Haemodialysis Department, Hald Ucad, Dakar, Senegal.

³Nephrology and Haemodialysis Department, CHR, St Louis, Gaston Berger University, Senegal.

⁴Nephrology and Haemodialysis Department, CHR, University of Thiès, Senegal.

yayuskanus@yahoo.fr

Corresponding Author: Dr Yaya Kane, Department of Health Science/Assane Seck University, Ziguinchor, Senegal.

Abstract

Mineral bone disorders (MBD) are almost constant complications in chronic hemodialysis patients. They cause an impairment in the quality of life and increase in cardiovascular morbidity and mortality. We are introducing the case of a 54-year-old woman, who was chronically on hemodialysis for 6 years, with a radial arteriovenous fistula as vascular access. Initially, the patient was reported having chronic tubulointerstitial nephritis. She was on hemodialysis three times a week. She developed secondary hyperparathyroidism. Clinically, she had diffuse arthralgias, bone pain mainly in the pelvis impeding walk; all this in a context of relative functional limitation of the lower limbs. As a result of paraclinical examination, serum calcium was 72 mg/l, phosphatemia was 42 mg/l. PTH returned to 2358 µg/ml and vitamin D tested using 25-OH-D was 20 mg/ml. Standard radiographs showed multiple geodes at the shoulder, lower extremity of the radius, trapezius, scaphoid, proximal phalanx head, spine and bilateral fracture lines of the femoral neck. Our patient was treated with calcium carbonate (e.g. Calcidia, in sachet), calcium-free phosphate binders (e.g. Renagel) and calcimimetics (e.g. Mimpara). Under medical treatment, there was a normalization of PTH and a decrease or even disappearance of the symptoms. This case shows that medical treatment for secondary hyperparathyroidism on hemodialysis patients, especially with calcimimetics, the use of which in our context is limited due to lack of availability.

Keywords: Secondary hyperparathyroidism, hemodialysis, calcimimetics, Ziguinchor, sub-Saharan Africa.

INTRODUCTION

In chronic hemodialysis (CHD), mineral and bone metabolism disorders (BMD) are early in chronic kidney disease, and progressively get worse over time [1]. Impaired phosphate excretion, reduced vitamin D activation, and compensatory increase in PTH secretion define secondary hyperparathyroidism, observable in most patients with advanced chronic renal failure [2]. KDIGO (Kidney Disease: improving Global Outcomes)

has recently suggested a more broadly defined term, i.e. “chronic kidney disease-mineral disorders” (CKD-MDB). [3]. Secondary hyperparathyroidism (PTH II) affected by hyperphosphoremia is part of renal osteodystrophy and is associated with an increased risk of cardiovascular mortality [4]. This clinical case highlights the difficulty of managing BMD in the situation of our developing country where biological and radiological explorations are expensive and not subsidized, in addition to the products used for

Secondary Hyper Parathyroidism, a Typical Mineral and Bone Disorder in a Chronic Hemodialysis Patient in Ziguinchor, Southern Senegal

treatment such as calcimimetics, hardly available on the local market. Our patient had the financial means that allowed her to take paraclinical examinations and get the therapeutic molecules, most often imported.

CLINICAL OBSERVATION

A 54-year-old woman received chronic kidney dialysis treatment at the hemodialysis center of the Ziguinchor Regional Hospital, 450 km away from Dakar, for 6 years, with a radial arteriovenous fistula for vascular access. Initially, the renal disease was supposed to be chronic tubulointerstitial nephritis. She was on a three sessions per week. She had mainly a story of heart disease with features of left heart failure. There was no previous aluminum toxicity. The diagnosis of hyperparathyroidism was established within 5 years after starting dialysis.

The blood pressure was stable (120/85 mmHg) when the clinical examination was carried out. There had no fever. She had arthralgia in her hips, knees, and right

shoulder. This was associated with widespread bone pain mainly at the level of the hip which restricted walk performance with the effect of relative periodical loss of limb function. The osteo-articular examination evidenced a limitation of movements and bone damage in the right wrist and pelvis. The biological assessment: serum calcium level was 82 mg/l (normal serum level between 85 and 105 mg/l), phosphatemia was 42 mg/l (normal level between 25 and 45 mg/l) The intact PTH 1-84 (parathyroid hormone) decreased to 2358 pg/ml (normal level between 6 and 50 pg/ml) and the 25 (OH) vitamin D concentration was 20 ng/ml (normal level is between 30 and 40 ng/ml). Hemoglobin was 11 g/dl erythropoietin 15,000 IU per 15 days. C reactive protein (CRP) was negative. Standard radiographs showed multiple bone cysts at the shoulder, lower extremity of radius, trapezoid, scaphoid, head of the thumb proximal phalanx and at the level of the spine, and bilateral stress fracture of the femoral necks (figure 1,2).



a. Cystic bone lesion at the lower end of the radius

b. Geode in the proximal phalanx of the thumb
Proximal phalanx geode of the thumb

c. Sub-periosteal resorption of distal phalanges

Fig 1. Standard radio of our patient's right hand, showing macrogeodes at the lower extremity of the radius, trapezoid, scaphoid and proximal phalanx of the dystrophic thumb.

Secondary Hyper Parathyroidism, a Typical Mineral and Bone Disorder in a Chronic Hemodialysis Patient in Ziguinchor, Southern Senegal



Fig 2. Standard radiographic view of a pathological bilateral fracture of the femoral necks in our patient.

The cervical ultrasound was eventless. Parathyroid scintigraphy was back to normal again.

Therapeutically, our patient was treated with calcium carbonate (Calcedia *: 1 sachet in the morning), a non-calcium-based phosphate binder sevelamer (Renagel * 800 mg: 1 tablet three times a day) and cinacalcet, a calcimimetic agent (Mimpara * 60 mg: 1 tablet twice daily and then dose adjustment according to the effects of PTH 1-84). In dialysis, she was submitted to high dialysate calcium concentration. Under medical treatment, there was a gradual decrease in PTH after 4-month monitoring. At 10 months of uninterrupted treatment, the biological parameters were returning to normal, with a decrease or even disappearance of the symptoms.

COMMENT

Secondary hyperparathyroidism can threaten life and functional prognosis in hemodialysis patients

as the results of cardiovascular and osteoarticular complications [5,6]. The prevalence of secondary hyperparathyroidism (HPT) is significant in developing countries, in the Ndiaye’s series with HPT in 45.5% of cases, in the different series of the literature as well, where the levels were between 30% and 60% on average. Studies by Llach et al [7], Sherrad et al. [8], Jebrane et al. [9], Seck et al. [10] and Benabdellah et al. [11] found a prevalence of 47.88%, 32.47%, 48.71%, 48% and 56.7% respectively (Table I).

High occurrence of the disease in our countries could easily be explained by the low socioeconomic level of our patients, who lack social coverage in most of cases. This makes a therapeutic treatment based on non-calcium phosphate binders and/or calcimimetics often difficult, or even unthinkable, given their high cost. Our patient had a high living standard which explained exceptional good care.

Table 1. Prevalence of HPT II in dialysis patients as part of the literature series

Authors	Years	Number of dialysis patients	HPT II	Prevalence
Llach et al. [7]	1986	142	68	47.88 %
Sherrard et al. [8]	1993	117	38	32.47 %
Ndiaye [12]	2008	40	14	45,5%
Guillaume et al. [13]	2010	502	158	30 %
Jebrane et al. [9]	2012	39	19	48.71 %
Seck et al. [10]	2012	118	57	48 %
Benabdellah et al. [11]	2013	83	47	56.7 %

Secondary Hyper Parathyroidism, a Typical Mineral and Bone Disorder in a Chronic Hemodialysis Patient in Ziguinchor, Southern Senegal

Our patient was 54 years old, which was consistent with the results found by Jebrane et al. (40.4 ± 10.6 years) [9], Montasser et al. (46 ± 15 years) [14], and Cherkaoui et al. (45 ± 10 years) [15]. In our case, HPT was diagnosed 5 years after onset of hemodialysis. This delay was similar to the finding by Montasser et al. (4 ± 2 years) [14] and Cherkaoui with 5 ± 4 years [15].

Traore et al. [16] found that 81.7% of patients were asymptomatic and 7.5% had bone pain and 5.3% muscle pain. However, in a recent series by Haddam et al. in Algeria (2015) [17], all patients had a clinical bone syndrome. Fibrous osteitis remains asymptomatic for a long time, bone pain usually occurs late. In our patient, we observed a high frequency of joint and bone pain, probably due to severity of the infections as well as the delay in these patients' care.

We found a very high levels of PTH, up to 2358 pg/ml (36 N), indicating a lack of adjustment and down-regulation in our work. These figures are significantly higher than those found in the Traoré et al. [16], Benabdellah et al. [11], Haddam et al. [17] series, with respectively 436.11 ± 200 µg/mL, 508 ± 380 µg/ml and 680 µg/ml of PTH on average.

The high parathyroid hormone (PTH) concentration in our patient could be explained by the fact that there are currently different kits for determining PTH level available on the market, with a wide range in the values obtained, as well as variation in defining hyperparathyroidism, depending on the series.

The dialysis reference values are 2 to 9 times the upper limit of the kit used (approximately 130-585 pg/ml for a 2nd generation assay).

Drug therapy for secondary hyperparathyroidism involves calcium, phosphate binders, calcitriol derivatives, and calcimimetics (cinacalcet Mimpara®). Our patient received the same type of treatment with a favorable evolution. In France, in the series by Guillaume et al., 51% of patients were treated with cinacalcet, of which 10% could be weaned after 12 months [13].

The modest use of calcimimetics in our center (in a patient), due to cost and lack of medical insurance in most of our patients, explains the fact that only our case had a control of hyperparathyroidism with PTH kept within the recommended targets for hemodialysis.

CONCLUSION

Secondary hyperparathyroidism is a common complication of chronic kidney failure. The care in sub-Saharan Africa is very restrictive, sometimes expensive. However, it still represents a challenge despite the recommendations of learned societies and knowledge of physiopathological mechanisms.

REFERENCES

- [1] Levin A, Bakris GL, Molitch M et al. Prevalence of abnormal serum vitamin D, PTH, calcium, and phosphorus in patients with chronic kidney disease: result of the study to evaluate early kidney disease. *Kidney Int*, 2007 ;71 :31.
- [2] Goodman WG, Quarles LD. Development and progression of secondary hyperthyroidism in chronic kidney disease: lessons from molecular genetics. *Kidney Int*, 2007 Jun 13 (Epub ahead of print).
- [3] Moe S, Drueke T, Cunningham J et al. Definition, evaluation, and classification of renal osteodystrophy: a position statement from kidney disease: improving global outcomes (KDIGO). *Kidney Int*, 2006 ;69 :1945.
- [4] Block GA, Hulbert-Shearon TE, Levin NW et al. Association of serum phosphorus and calcium X phosphate product with mortality risk in chronic hemodialysis patients: a national study. *Am J Kidney Dis* 1998 ;31 :607-17.
- [5] Block G, Klassen P, Kim J. Relationship between serum calcium and mortality risk in hemodialysis patient. *Am J kidney dis* 2003 ;41: A16.
- [6] Block GA, Klassen P, Lazarus JM, Ofsthun N, Lowrie EG, Chertow GM. Mineral metabolism, mortality and morbidity in maintenance hemodialysis. *J Am soc nephrol* 2004 ;15: 2208-18.
- [7] Llach F, Felsenfeld A, Coleman M, et al. The natural course of dialysis osteomalacia. *Kidney Int* 1986; 29 :74-9.
- [8] Sherrard D, Hercz G, Pei Y, et al. The spectrum of bone disease in end-stage renal failure: An evolving disorder. *Kidney Int* 1993 : 43 :436-42.
- [9] **Jabrane M.** Les troubles minéraux et osseux chez les hémodialysés au service de Néphrologie-

Secondary Hyper Parathyroidism, a Typical Mineral and Bone Disorder in a Chronic Hemodialysis Patient in Ziguinchor, Southern Senegal

- Hémodialyse du CHU Med VI de Marrakech. Thèse Med. Marrakech, N° 129, 2012.
- [10] Seck SM, Dahaba M, Ka EF et al. Mineral and bone disease in black african hemodialysis patients: a report from senegal. *Nephrourol Mon.* 2012 Fall; 4(4):613-6.
- [11] **Benabdellah N, Karimi I, Bentata Y, et al.** Statut phospho-calcique en hémodialyse chronique dans l'Oriental Marocain: évaluation de l'adhésion aux recommandations K/DOQI et KDIGO. *Pan Afr Med J.* 2013; 16: 23.
- [12] Ndiaye N. L'ostéodystrophie rénale : Aspects épidémiologiques, cliniques et thérapeutiques au centre d'hémodialyse du CHU Le Dantec. Thèse Med, Dakar, N° 65, 2008.
- [13] Guillaume Jean. Améliorer la prise en charge de l'hyperparathyroïdie secondaire au Maghreb : pour faire enfin disparaître ces tumeurs brunes. *Nephrol Ther* ; Janvier 2016, 832.
- [14] Montasser D, Benyahia M, Zajjari Y et al. Traitement chirurgical de l'hyper parathyroïdie secondaire chez l'hémodialysé chronique. *Nephrol Ther* 2011; 7 (5): 314.
- [15] Cherkaoui A, G. El Badaoui, G. El Miftah M et al. Profil de l'hyperparathyroïdie secondaire en dialyse. *Nephrol Ther* 2013; 9: 5-311.
- [16] Traoré D, Traoré B, Nientao I et al. Etude épidémioclinique de l'hyperparathyroïdie secondaire à l'insuffisance rénale chronique dans le service de néphrologie et d'hémodialyse du CHU du point G. *Ann Endocrinol* 2015 ; 76 (4) :479-480.
- [17] Haddam A.E.M, Fedala N. S, Chentli F, D. Meskine. L'hyperparathyroïdie secondaire à l'insuffisance rénale : à propos d'une série de 15 patients. *Ann Endocrinol* 2015; 76: 476-477.

Citation: Kane Yaya, Bangoura Mohamed, Dia Gueye Diatou, et al. *Secondary Hyper Parathyroidism, a Typical Mineral and Bone Disorder in a Chronic Hemodialysis Patient in Ziguinchor, Southern Senegal. Archives of Nephrology.* 2018; 2(1): 01-05.

Copyright: © 2018 Kane Yaya, Bangoura Mohamed, Dia Gueye Diatou, et al. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*